arXiv:1604.06012v1 [astro-ph.SR] 20 Apr 2016

A note on cement in asteroids

G. Bilalbegović

Department of Physics, Faculty of Science, University of Zagreb, Bijenička 32, 10000 Zagreb, Croatia

Abstract

Cement mineral tobermorite was formed in hydrothermal experiments on alternation of calcium-aluminum-rich inclusions (CAIs) in carbonaceous chondrite meteorites. Unidentified bands at 14 μm were measured for CAIs and the matrix of the Allende meteorite sample, as well as for Hektor and Agamemnon asteroids. The presence of cement nanoparticles may explain the feature at 14 μm.

Keywords: Asteroids, composition; Infrared observations; Meteorites; Asteroids, surfaces; Trojan asteroids

Cement is one of the most used materials on the Earth. Concrete is made from cement paste, sand, and stone. Cement paste consists of cement powder and water. The main component of cement, calcium-silicate-hydrate, has a granular complex structure at nanoscales (see Fig. 1 in Allen et al. (2007)). Although concrete and cement we use are man-made materials, several cement crystals are naturally formed on the Earth. The structure of cement is sometimes described using minerals tobermorite and jennite, but many other natural terrestrial forms of calcium-silicate-hydrates exist (see Table 1 in Richardson (2008)). For example, even tobermorite is not the unique crystal, but appears in three main forms, known as 9 Å, 11 Å, and 14 Å tobermorites. These systems vary in the interlayer distance. It is found that more than 30 stable phases of calcium-silicate-hydrates form in hydrothermal environments (Shaw et al., 2000). Two of these phases are 11 Å tobermorite ($Ca_5Si_6O_{16}(OH)_2 \cdot 4\ H_2O$) and xonotlite ($Ca_6Si_6O_{17}(OH)_2$).

A recent study of cement nanoparticles revealed that, together with typ-

Email address: goranka.bilalbegovic@gmail.com (G. Bilalbegović)

ical bands of amorphous silicates at 10 and 18 μm, these structures show specific infrared (IR) features at 14 μm (Bilalbegović et al., 2014). Figs. 2-4 in Bilalbegović et al. (2014) show that bands of cement nanoparticles exist in near-, mid- and far-IR spectral regions. IR bands are described and features with higher intensities are listed in Table 1 of Bilalbegović et al. (2014). Measurements of IR spectra for the cement paste showed that the feature at 14 μm exists for a range of compositions and it is generated by the Si-O-Si bending vibrations (Garbev et al., 2007). However, it is known that the frequency of these bending vibrations changes with silicate structure and polymerization. It should be expected that exact positions of bands close to 14 μm vary with the composition and size of cement nanoparticles.

It is possible that cement forms in various environments in Space. The Infrared Space Observatory (ISO) and the Spitzer telescope observed unidentified IR features at 14 μm for several stars: HD 44179 (Red rectangle), HD 45677, and IRC+10420 (Molster et al., 2002; Markwick-Kemper et al., 2005). Recent data for the planetary nebula BD+303639 taken by the FORCAST instrument on board of the Sofia telescope have also shown the unidentified 14 μm band (Guzman-Ramirez, personal communications). Table 1 compares positions of IR bands at 14 μm, which were calculated, measured, or observed for several objects on the Earth and in Space.

It is known that some meteorites contain calcium-aluminum-rich inclusions (CAIs) that are either pre-solar grains, or the first material condensed in the Solar nebula (Krot et al., 2005). Nomura and Miyamoto (1998) carried out experiments on alternation of CAIs in carbonaceous chondrites to study aqueous alternation in parent asteroids. They investigated some of the oldest phases in the Solar System: gehlenite, spinel, and diopside, as well as mixtures of gehlenite with SiO_2 and Al_2O_3. One of the products obtained in hydrothermal experiments from the mixture of gehlenite with SiO_2 was tobermorite. This mineral is one of Ca-Si hydrates and exists as natural cement on the Earth (Richardson, 2008). Nomura and Miyamoto (1998) suggested that the formation of Ca-Si hydrates started when gehlenite was disolved by a fluid and Ca, Al, and Si atoms were released. Tobermorite crystallized when sufficient numbers of Ca and Si atoms were present. Nomura and Miyamoto (1998) wrote that Shoji, T. (1974a,b) in hydrothermal experiments obtained another Ca-Si hydrate mineral xonotlite. The hydrothermal formation of xonotlite was later investigated in details (Shaw et al., 2000). Hydrous minerals produced in asteroids by the process of aqueous alternation could convert into anhydrous ones by thermal metamorphism. Nomura and Miyamoto

(1998) found that tobermorite exists in a whisker form and converts to wollastonite with increasing temperature. It is possible that both tobermorite and wollastonite exist in the same environment if thermal metamorphism is not complete.

The Allende meteorite is from the oxidized subgroup of the CV carbonaceous chondrites. Allende was affected by aqueous alternation on the parent body (Krot et al., 1998). It is known that both gehlenite (Nomura and Miyamoto, 1998) and wollastonite (Miyamoto et al., 1979; Krot et al., 1998) are present in Allende. Therefore, it is possible that tobermorite and other calcium-silicate-hydrates were also formed in the parent asteroid body and could be detected in Allende. CAIs are minor components of chondrite meteorites. In CV chondrites refractory inclusions (CAIs and AOAs (amoeboid olivine aggregates) together) make about 10 vol. 10% (Scott and Krot, 2005). Therefore, the amount of cement expected to form in the processes suggested by Nomura and Miyamoto (1998) is not big. Morlok and coworkers studied mid-IR spectra (between 2.5 μm and 16 μm) of CAIs separated from Allende (Morlok et al., 2008). The sample analyzed was from the collection of the Natural History Museum in London and it was collected immediately after its fall. Therefore, this sample was not substantially changed by conditions on the Earth. The sample measured was a fine-grained, submicron-sized powder. It was difficult to characterize the minerals. Therefore, mixtures of several phases were analyzed. CaO i SiO_2 were detected. Table 2 shows IR bands at 14 μm measured for materials in the sample of Allende (Morlok et al., 2008). Morlok and coworkers suggested that a carrier of the broad feature between 14.0 μm and 15.0 μm could be spinel. For example, Chihara et al. (2000) studied the optical constants of three crystalline spinels ($MgAl_2O_4$) in the mid- and far-infrared spectral regions. Two samples were natural spinels (white and pink), whereas one was synthesized. The pink spinel included a small amount of Cr, and synthetic spinel was Al_2O_3 rich. Bands at 14 μm were measured for the pink spinel (14.7 μm in the absorption spectrum), the white spinel (14.8 μm in absorption), and the synthetic spinel (14.6 μm in absorption and 14.0 μm in reflection). Morlok and coworkers proposed that a position within the broad band probably depends on the spinel's iron content. However, IR spectra of iron-bearing spinels were measured (Cloutis et al., 2004; Jackson et al., 2014). Cloutis and coworkers found that the best correlation exists between Fe^{2+} content and wavelength positions at 0.46, 0.93, 2.8, 12.3, 16.2, and 17.5 μm. For Fe^{3+} content the best correlation is at 0.93 μm. The authors did not mention any relevance of the (14-15) μm

spectral region for a broad range of chemical compositions of iron-bearing spinels they studied (Cloutis et al., 2004; Jackson et al., 2014). IR features of cement can explain bands at 14 μm measured by Morlok and coworkers.

Emery et al. (2006) analyzed spectra of Trojan asteroids 624 Hektor, 911 Agamemnon, and 1172 Aneas, obtained by the IR spectrograph (IRS) on the Spitzer Space Telescope. After comparisons of their data with available spectral libraries for meteorites and minerals, these authors concluded that small silicate grains suspended in a transparent matrix are the best candidates to reproduce mid-IR spectral shapes of asteroids they studied, such as the Si-O stretch fundamental at 10 μm and the Si-O bend fundamental at $\sim$18 to 25 μm. However, comparisons with existing spectral databases for meteorites and minerals did not explain all details of asteroid spectra. A small, but visible, spectral feature between 14 μm and 15 μm was found for Hektor and Agamemnon. Cement nanoparticles are good candidates for carriers of this feature.

Hydrated minerals in asteroids are traced by absorption features at 0.7 and 3 μm. Observations of Trojan asteroids in the visible (Bendjoya et al., 2004) and near-IR (Cruikshank et al., 2001; Emery and Brown, 2003) spectral regions did not show these bands. Therefore, till now there is no evidence of hydrated minerals on surfaces of Trojan asteroids and a route to the cement formation on Hektor and Agamemnon based on altered CAIs is not supported by available observations. The Tagish lake carbonaceous chondrite shows similar spectra as the D-type asteroids and it was proposed as their analog (Hiroi et al., 2001). A recent analysis shows that possible parent bodies for Tagish Lake are very rare in the group of Jupiter Trojans (Vernazza et al., 2013). The Tagish Lake meteorite is carbon rich and has high concentrations of presolar grains, as well as carbonate minerals. The presence of these ingredients is consistent with the present understanding of compositions of the D-type asteroids. However, Trojans have redder spectral slopes and the Tagish lake meteorite is aqueously altered. Several ideas have been put forward to explain the non-detection of hydrated minerals on Trojans. Cruikshank et al. (2001) proposed that hydrated minerals on the surface of Hektor could be masked with low-albedo materials, such as carbon. They also suggested that the strength of 3 μm band decreases with heliocentric distance because of increasing abundance of carbon. Dehydration by micrometeorite bombardments of the surface of some D-type asteroids was also proposed (Hiroi et al., 2001). Future high-resolution observations of Trojan asteroids, as well as a fall of a meteorite similar to the Jupiter

Trojan D-type asteroids, would be able to settle this problem.

Experiments on alternation of CAIs by Nomura and Miyamoto (1998) shown that cement forms in carbonaceous chondrites. Ca atoms are supplied by gehlenite in this hydrothermal scenario of cement formation. It is known that gehlenite (Al-rich melilite) is a typical mineral in CAIs (Nomura and Miyamoto, 1998; Morlok et al., 2008). Emery et al. (2006) used a modified standard thermal model to estimate the surface temperatures of Trojan asteroids they studied. Because of insufficient information on the physical properties, they also varied several parameters to find the best fit to the measured spectral energy distribution and found that the Hector's surface temperature is 179.4 K. Nomura and Miyamoto (1998) performed their experiments on alternation of CAIs in carbonaceous chondrites where tobermorite was formed at 473.15 K. The presence of water ice and other ices in Trojan asteroids was studied (Cruikshank et al., 2001; Yang and Jewitt, 2007; Guilbert-Lepoutre, 2014). Present understanding is that Trojans have a large fraction of water ice covered by a refractory mantle (Emery et al., 2015). Yang and co-workers suggested that in the young Solar system short-lived radionuclides heated Trojans and their ices melted (Yang et al., 2013). A mechanical mixture of water, gehlenite and other primitive silicates, as well as other soluble materials circulated. These conditions were on some asteroids favorable for the formation of cement nanoparticles.

Cement, as a specific silicate, should be included in spectral databases for future studies of asteroids and meteorites.

This work is supported by the HRZZ Research Project "Stars and dust: structure, composition and interaction" and the QuantiXLie Center of Excellence. I would like to thank Lizette Guzman-Ramirez for interesting discussions about the BD+303639 planetary nebula and IR spectra. This research has made use of NASA's Astrophysics Data System Bibliographic Services.

References

Allen, A. J., Thomas, J. J., Jennings, H. M., 2007. Composition and density of nanoscale calcium-silicate-hydrate in cement. Nature Mater. 6, 311.

Table 1: IR bands at 14 μm for several cement structures and corresponding unidentified features in astrophysical objects. Data are from: (a) Bilalbegović et al. (2014), (b) Garbev et al. (2007), (c) Markwick-Kemper et al. (2005), (d) Molster et al. (2002), (e) Emery et al. (2006).

Source	State	IR at 14 μm
$Ca_4Si_4O_{14}H_4$	nanoparticle, crystalline precursor	14.44, 14.55 [(a)]
$Ca_6Si_3O_{13}H_2$	nanoparticle, amorphous	14.10 [(a)]
$Ca_{12}Si_6O_{26}H_4$	nanoparticle, amorphous	14.58 [(a)]
Cement paste	compositions $CaO/SiO_2 = 0.2$ -1.5	14.88 - 14.97 [(b)]
Red Rectangle (Spitzer)	unknown	14.-16. [(c)]
Red Rectangle (ISO)	unknown	14.19 [(d)]
HD 45677	unknown	14.3 [(d)]
IRC+10420	unknown	14.15 [(d)]
Hektor, Agamemnon	unknown	14.-15. [(e)]

Table 2: IR bands at 14 μm for the matrix and CAIs of the Allende meteorite sample (Morlok et al., 2008).

Object	Bands at 14 μm
Matrix	14.29, 14.97
RIM 1	14.62, 14.88
RIM 2	14.62, 14.88
Interior 1	14.12, 14.39, 14.97
Interior 2	14.21, 14.88
Interior 4	14.88
Interior (close RIM)	14.97

Bendjoya, P., Cellino, A., Di Martino, M., Saba, L., 2004. Spectroscopic observations of Jupiter Trojans. Icarus 168, 374–384.

Bilalbegović, G., Maksimović, A., Mohaček-Grošev, V., Aug. 2014. Do cement nanoparticles exist in space? Mon. Not. R. Astron. Soc. 442, 1319.

Chihara, H., Tsuchiyama, A., Koike, C., Sogawa, H., 2000. The Optical Constant of Crystaline Spinel. In: Garzón, G., Eiroa, C., de Winter, D., Mahoney, T. J. (Eds.), Disks, Planetesimals, and Planets. Vol. 219 of ASP Conf. Ser. p. 150.

Cloutis, E. A., Sunshine, J. M., Morris, R. V., 2004. Spectral reflectance-compositional properties of spinels and chromites: Implications for planetary remote sensing and geothermometry. Meteorit. Planet. Sci. 39, 545.

Cruikshank, D. P., Dalle Ore, C. M., Roush, T. L., Geballe, T. R., Owen, T. C., de Bergh, C., Cash, M. D., Hartmann, W. K., 2001. Constraints on the composition of Trojan asteroid 624 Hektor. Icarus 153, 348.

Emery, J. P., Brown, R. H., 2003. Constraints on the surface composition of Trojan asteroids from near-infrared (0.8-4.0 μm) spectroscopy. Icarus 164, 104–121.

Emery, J. P., Cruikshank, D. P., Van Cleve, J., 2006. Thermal emission spectroscopy (5.2-38 μm) of three Trojan asteroids with the Spitzer Space Telescope: Detection of fine-grained silicates. Icarus 182, 496.

Emery, J. P., Marzari, F., Morbidelli, A., French, L. M., Grav, T., 2015. The complex history of Trojan asteroids. In: Michel, P., DeMeo, F. E., Bottke, W. F. (Eds.), Asteroids IV. University of Arizona Press, Tucson, AZ, p. 203.

Garbev, K., Stemmermann, P., Black, L., Breen, C., Yarwood, J., Gasharova, B., 2007. Structural features of CSH(I) and its carbonation in air - a Raman spectroscopic study. Part I: fresh phases. J. Am. Ceram. Soc. 90, 900.

Guilbert-Lepoutre, A., 2014. Survival of water ice in Jupiter Trojans. Icarus 231, 232.

Hiroi, T., Zolensky, M. E., Pieters, C. M., 2001. The Tagish Lake Meteorite: A Possible Sample from a D-Type Asteroid. Science 293, 2234–2236.

Jackson, C. R. M., Cheek, L. C., Williams, K. B., Hanna, K. D., Pieters, C. M., Parman, S. W., Cooper, R. F., Dyar, M. D., Nelms, M., Salvatore, M. R., 2014. Visible-infrared spectral properties of iron-bearing aluminate spinel under lunar-like redox conditions. Am. Mineral. 99, 1821.

Krot, A. N., Petaev, M. I., Scott, E. R. D., Choi, B.-G., Zolensky, M. E., Keil, K., 1998. Progressive alteration in CV3 chondrites: More evidence for asteroidal alteration. Meteorit. Planet. Sci. 33, 1065.

Krot, A. N., Yurimoto, H., Hutcheon, I. D., MacPherson, G. J., 2005. Chronology of the early Solar System from chondrule-bearing calcium-aluminium-rich inclusions. Nature 434, 998.

Markwick-Kemper, F., Green, J. D., Peeters, E., 2005. Spitzer detections of new dust components in the outflow of the Red rectangle. Astrophys. J. 628, L119.

Miyamoto, M., Onuma, N., Takeda, H., 1979. Wollastonite whiskers in the Allende meteorite and their bearing on a possible post-condensation process. Geochem. J. 13, 1.

Molster, F. J., Waters, L. B. F. M., Tielens, A. G. G. M., Barlow, M. J., 2002. Crystalline silicate dust around evolved stars. I. The sample stars. Astron. Astrophys. 382, 184.

Morlok, A., Köhler, M., Grady, M. M., 2008. Mid-infrared spectroscopy of refractory inclusions (CAIs) in CV and CO chondrites. Meteorit. Planet. Sci. 43, 1147.

Nomura, K., Miyamoto, M., 1998. Hydrothermal experiments on alteration of Ca-Al-rich inclusions (CAIs) in carbonaceous chondrites: implication for aqueous alteration in parent asteroids. Geochim. Cosmochim. Ac. 62, 3575.

Richardson, I. G., 2008. The calcium silicate hydrates. Cem. Concr. Res. 38, 137.

Scott, E. R. D., Krot, A. N., 2005. Chondritic Meteorites and the High-Temperature Nebular Origins of Their Components. In: Krot, A. N., Scott, E. R. D., Reipurth, B. (Eds.), Chondrites and the Protoplanetary Disk. Vol. 341 of ASP Conf. Ser. p. 15.

Shaw, S., Clark, S. M., Henderson, C. M. B., 2000. Hydrothermal formation of the calcium silicate hydrates, tobermorite ($Ca_5Si_6O_{16}(OH)_2 \cdot 4\,H_2O$) and xonotlite ($Ca_6Si_6O_{17}(OH)_2$): an in situ synchrotron study. Chem. Geol. 167, 129.

Shoji, T., 1974a. $Ca_3Al_2(SiO_4)_3$-$Ca_3Si_2(O_4H_4)_3$ Series Garnet: Composition and Stability. J. Mineral. Soc. Japan 11, 359, in Japanese.

Shoji, T., 1974b. Phase relations in the System CaO-Al_2O_3-SiO_2-H_2O. J. Mineral. Soc. Japan 12, 1, in Japanese.

Vernazza, P., Fulvio, D., Brunetto, R., Emery, J. P., Dukes, C. A., Cipriani, F., Witasse, O., Schaible, M. J., Zanda, B., Strazzulla, G., Baragiola, R. A., 2013. Paucity of Tagish Lake-like parent bodies in the Asteroid Belt and among Jupiter Trojans. Icarus 225, 517–525.

Yang, B., Jewitt, D., 2007. Spectroscopic search for water ice on Jovian Trojan asteroids. Astron. J. 134, 223.

Yang, B., Lucey, P., Glotch, T., 2013. Are large Trojan asteroids salty? An observational, theoretical, and experimental study. Icarus 223, 359.